

NOTICE: When government or other drawings, specifications or other data are used for any purpose other than in connection with a definitely related government procurement operation, the U. S. Government thereby incurs no responsibility, nor any obligation whatsoever; and the fact that the Government may have formulated, furnished, or in any way supplied the said drawings, specifications, or other data is not to be regarded by implication or otherwise as in any manner licensing the holder or any other person or corporation, or conveying any rights or permission to manufacture, use or sell any patented invention that may in any way be related thereto.

AD No. 453813

DDC FILE COPY.


4 5 3 8 1 3

Reprinted from:

TETRAHEDRON LETTERS

*The International Organ for the Rapid Publication
of Preliminary Communications in Organic Chemistry*

REPRODUCTION IN WHOLE OR
IN PART IS PERMITTED FOR ANY
PURPOSE OF THE U.S. GOVERNMENT.



PERGAMON PRESS

OXFORD • LONDON • NEW YORK • PARIS

(6) INVERSION RATE IN A PRIMARY GRIGNARD REAGENT

Gideon Fraenkel, Don T. Dix and David G. Adams
Department of Chemistry, The Ohio State University, Columbus 10, Ohio

(Received 27 July 1964; in revised form 31 August 1964)

There is a wealth of chemical evidence concerning carbanion inversion phenomena in organometallic reagents. ~~(1-6)~~

We now report a direct physical measurement of the inversion rate in a primary Grignard reagent using the techniques of NMR ~~(7,8,9)~~

The Grignard reagent in ether was prepared from 2-methyl-butyl bromide. (10) The NMR absorption at 60 Mc and -30° of the $\text{CH}_2\text{-Mg}\cdots\text{hydrogens}$ in the Grignard reagent is illustrated in Fig. 1a; it consists of the AB portion of the familiar ABX system (11) where $J_{AB} = \pm 12.4$, $J_{AX} = +7.65$ and $J_{BX} = +4.35$ cps respectively and $\delta_A - \delta_B = 0.170$ ppm. The methylene hydrogens are magnetically non-equivalent because H_A is always in an environment different from that around H_B regardless as to the rate of rotation about the $\text{CH}_2\text{-C}$ (asymmetric) bond. (12-14) Only by exchanging places simultaneously can H_A and H_B become magnetically equivalent. Such an effect could be accomplished by inversion at the Mg-CH_2 - carbon atom. (15) This is indeed the case for with increasing temperature the $\text{CH}_2\text{-Mg}$ multiplet in the Grignard NMR spectrum collapses to a doublet, A_2X (16) by 70° . See Fig. 1a - e.

3157




FIG. 1

NMR absorption of CH_2Mg multiplet in 2-methyl-butylmagnesium bromide, 1M, a, -30° ; b, -5° ; c, 21.5° ; d, 40° ; e, 70° .

We have calculated inversion frequencies from the line-shapes of the collapsing $\text{CH}_2\text{-Mg}$ multiplet. At -30° the latter consists of two overlapping AB quartets generated by the spin-states of H_X , $-\text{CH}_X\text{CH}_2\text{Mg}$. They are lines 1,3,4, and 7 and lines 2,5,6, and 8 in Fig. 1a. The distance between the quartet centers is $\frac{1}{2}(J_{AX} + J_{BX})$. It is assumed that during an inversion the spin states of the methylene hydrogens do not change. Hence with rapid inversion the quartets collapse independently to single lines at their center. (17) Using Saunders' computer program for determining the line-shape of an AB system undergoing $\text{H}_A \rightleftharpoons \text{H}_B$ exchange at different rates (18) we have superposed the line-shapes for the two collapsing quartets described above and compared the calculated spectra with the observed results. At 25° the mean lifetimes between inversions is 2.4×10^{-2} sec, ΔH^\ddagger and ΔS^\ddagger are 12 ± 1 kcal and

-17 ± 3 eu. *Y. P. Tang, J. Am. Chem. Soc. 81, 5155 (1959)*
 We have obtained very similar results for 2-methyl-butyllithium in pentane.

These results indicate that primary Grignard reagents and organo-lithium compounds undergo rapid inversion at ordinary temperatures.

The kinetics of these processes will be reported later. Elucidation of the mechanisms of these inversion processes must await completion of these kinetic studies.

Acknowledgement This research was generously supported by the Petroleum Research Fund of the American Chemical Society and the U. S. Air Force Office of Scientific Research.

REFERENCES

1. M. S. Kharasch and O. Reinuth, "Grignard Reactions of Organometallic Substances", Prentice Hall, New York, 1954.
2. R. E. Dessey and B. Jones., J. Org. Chem., 24, 1685 (1959).
3. D. J. Cram et al. J. Amer. Chem. Soc., 81, 5740-5750 (1959).
4. R. C. Letsinger, ibid., 72, 4842 (1950).
5. H. M. Walborsky and F. J. Impasato, ibid., 81, 5835 (1959).
6. H. M. Walborsky and A. E. Young, Abstracts of Papers, Organic Division, American Chemical Society Meeting, March 1962.
7. J. A. Pople, W. G. Schneider and H. J. Bernstein, High Resolution Nuclear Magnetic Resonance, McGraw-Hill Book Co., Inc. New York, 1959, Chap. 10.
8. G. Fraenkel, D. G. Adams and J. Williams Tetrahedron Letters, 161 (1963).
9. S. Alexander, J. Chem. Phys., 37, 966 (1962).
10. Obtained by the method of Crombie: L. Crombie and G. H. Harper, J. Chem. Soc., 2688 (1950).
11. Ref. 7 Chap. 6.
12. P. M. Nair and J. D. Roberts, J. Amer. Chem. Soc., 79, 4565 (1957).
13. H. S. Gutowsky, J. Chem. Phys., 37, 2196 (1962).
14. J. A. Pople, Mol. Phys., 1, 1 (1958).
15. M. Saunders and F. Yamada, J. Amer. Chem. Soc., 85, 1882 (1964).
16. Roberts has attributed the temperature dependence of the NMR spectra of certain Grignard reagents to different inversion rates but has not reported the inversion rates, G. M. Whitsides, F. Kaplan and J. D. Roberts, J. Amer. Chem. Soc., 85, 6217 (1963).
17. A similar method was used to obtain the rate of rotation in N,N-dimethylformamide, G. Fraenkel and C. Franconi, ibid., 82, 4478 (1960).
18. We thank Professor Martin Saunders, Yale University for giving us a copy of his program.